

HYPOTONIA – FLOPPY BABY

Supporting information

This guideline has been prepared with reference to the following:

Dyke M and Jackson N. Joint Trust Guideline for Assessment of a Floppy Baby (Neonatal Hypotonia). 2016. James Paget University Hospitals NHS Foundation Trust and Norfolk and Norwich University Hospitals NHS Foundation Trust

Bodamer O and Miller G. Approach to the infant with hypotonia and weakness. In: UpToDate. 2016

Auckland District Health Board. Neonatal Hypotonia. 2015

Lisi EC and Cohn RD. Genetic evaluation of the pediatric patient with hypotonia: perspective from a hypotonia specialty clinic and review of the literature. *Dev Med Child Neurol.* 2011;53:586-99

<http://onlinelibrary.wiley.com/doi/10.1111/j.1469-8749.2011.03918.x/full>

Children's Hospital Orange County. Neonatal Hypotonia - Clinical Approach to Floppy Baby. 2011. CHOC Children's

<https://specialists.chocchildrens.org/wp-content/uploads/2011/08/Neonatal-Hypotonia.pdf>

Leyenaar J, Camfield P, Camfield C. A schematic approach to hypotonia in infancy. *Paediatr Child Health.* 2005;10:397-400

<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2722561/>

van Toorn R. Clinical approach to the floppy child. *CME.* 2004;22:449-55

<http://www.ajol.info/index.php/cme/article/viewFile/44002/27518>

Last amended October 2016
Last reviewed December 2021